

# NeuroMed7<sup>®</sup> Application Protocol

Proper protocol for application of NeuroMed7 is extremely important for positive outcome effect:

- **Use approximately a “US quarter dollar” size amount per 4”x4” area immediately prior to procedure**
- **Avoid use on large open wounds, broken skin, eyes, ears, in the nostrils and mouth**
- This volume of product is sufficient for the surface area of, for example, half of a face or an average size adult male shoulder
- Depending on the skin type and surface area (for example, an entire face), you may use approximately double that amount
- We recommend using no more than 3 full size doses at one time (or about 10% of surface area of skin)
- **With a gloved hand, evenly distribute the cream and rub vigorously in a circular motion for at least 30 seconds**
- This is a “desensitizing” or comfort increasing agent, and you should see an initial onset of action in 5 – 10 minutes
- The peak action begins in approximately 15 – 20 minutes
- For procedures lasting longer than 60 minutes, reapply as described above, within the parameters also described above
- For smaller surface areas (such as the eyebrow area) a slightly smaller amount can be used
- For more aggressive or deeper procedures some clinicians use an occlusive dressing for an improved efficacy – the occlusive dressing has been applied for the full 15 – 20 minutes prior to procedure – occlusion should be determined by a licensed medical practitioner, taking into account potential toxicity risk factors
- NeuroMed7<sup>®</sup> can be applied in more liberal volume where applicable as deemed necessary by a licensed clinician
- Physician should take into account total topical anesthetic dosage application volume if using this product in combination with other anesthetic products

For more information contact your local  
Representative or Sambria Pharmaceuticals:  
info@sambriapharma.com  
www.sambriapharma.com  
888.246.6601

